arXiv:1411.3037v2 [math.DG] 30 Jan 2015

THE GAUSS MAP AND TOTAL CURVATURE OF COMPLETE MINIMAL LAGRANGIAN SURFACES IN THE COMPLEX TWO-SPACE

REIKO AIYAMA, KAZUO AKUTAGAWA, AND YU KAWAKAMI

Abstract. The purpose of this paper is to reveal the relationship between the total curvature and the global behavior of the Gauss map of a complete minimal Lagrangian surface in the complex two-space. To achieve this purpose, we show the precise maximal number of exceptional values of the Gauss map for a complete minimal Lagrangian surface with finite total curvature in the complex two-space. Moreover, we prove that if the Gauss map of a complete minimal Lagrangian surface which is not a Lagrangian plane omits three values, then it takes all other values infinitely many times.

1. Introduction

There are many similarities between surfaces in Euclidean 3-space $\mathbf{R}^3$ and Lagrangian surfaces in the complex 2-space $\mathbf{C}^2$, in particular, the case of minimal surfaces. In fact, there exists a representation for a minimal Lagrangian surface $M(\subset \mathbf{C}^2)$ in terms of holomorphic data, similar to Weierstrass representation for a minimal surface in $\mathbf{R}^3$ (cf. [26]). Moreover, the Gauss map g of M is a holomorphic map to the unit 2-sphere S^2. On the representation for M, Chen-Morvan [5] proved that there exists an explicit correspondence in $\mathbf{C}^2$ between minimal Lagrangian surfaces and holomorphic curves with a nondegenerate condition. Indeed, this correspondence is given by exchanging the orthogonal complex structure J in $\mathbf{C}^2$ to another one on $\mathbf{R}^4 = \mathbf{C}^2$. For the complete case, this result can also be proved from [22, Theorem II] and the well-known fact [10] that any minimal Lagrangian submanifold in $\mathbf{C}^n$ is stable. More generally, Hélein-Romon [11, 12] and the first author [1, 2] proved that every Lagrangian surface S in $\mathbf{C}^2$, not necessarily minimal, is represented in terms of a plus spinor (or a minus spinor) of the spin$^{\mathbf{C}}$ bundle $(\underline{\mathbf{C}}_S \oplus \underline{\mathbf{C}}_S) \oplus (K_S^{-1} \oplus K_S)$ satisfying the Dirac equation with potential (see [2, Section 1] for details). Here, $\underline{\mathbf{C}}_S$ and K_S denote respectively the trivial complex line bundle and the canonical complex line bundle of S. Remark that the representation in terms of plus spinors in $\Gamma(\underline{\mathbf{C}}_S \oplus \underline{\mathbf{C}}_S) = \Gamma(S \times \mathbf{C}^2)$ given by the first author is a natural generalization of the one given by Chen-Morvan. Combining these results, we get the following:

2010 *Mathematics Subject Classification.* Primary 53C42, 53D10; Secondary 30D35, 53D12.

Key words and phrases. Lagrangian surface, Gauss map, total curvature, exceptional value.

The second author is supported in part by the Grant-in-Aid for Challenging Exploratory Research, Japan Society for the Promotion of Science, No. 24654009.

The third author is supported by the Grant-in-Aid for Young Scientists (B), Japan Society for the Promotion of Science, No. 24740044.

THEOREM 1.1. ([5], [1, 2]) *Let M be a Riemann surface with an isothermal coordinate $z = u+\mathbf{i}\,v$ around each point. Let $F = (F_1, F_2)\colon M \to \mathbf{C}^2$ be a holomorphic map satisfying $|S_1|^2 + |S_2|^2 \neq 0$ everywhere on M, where $S_1 := (F_2)'_z = dF_2/dz$ and $S_2 := -(F_1)'_z = -dF_1/dz$. Then*

$$f = \frac{1}{\sqrt{2}} e^{\mathbf{i}\,\beta/2}(F_1 - \mathbf{i}\,\overline{F_2}, F_2 + \mathbf{i}\,\overline{F_1}) \tag{1}$$

is a minimal Lagrangian conformal immersion from M to $\mathbf{C}^2$ with constant Lagrangian angle $\beta \in \mathbf{R}/2\pi\mathbf{Z}$. The induced metric ds^2 on M by f and its Gaussian curvature K_{ds^2} are respectively given by

$$ds^2 = (|S_1|^2 + |S_2|^2)|dz|^2, \qquad K_{ds^2} = -2\frac{|S_1(S_2)_z - S_2(S_1)_z|}{(|S_1|^2 + |S_2|^2)^3}. \tag{2}$$

The Gauss map g of M is also given by

$$g = [-S_2 : S_1] = (-S_2/S_1)\colon M \to \mathbf{C}P^1 = \mathbf{C} \cup \{\infty\} \simeq S^2. \tag{3}$$

Here, by the identification of S^2 with $S^2(1)\times\{(e^{\mathbf{i}\beta}, 0)\} \subset \mathbf{R}^3\times\mathbf{R}^3$, g can be regarded as the generalized Gauss map of $F(M)$ in $\mathbf{R}^4 = \mathbf{C}^2$ (cf. [13], [14]). *Conversely, every minimal Lagrangian immersion $f\colon M \to \mathbf{C}^2$ with constant Lagrangian angle β is congruent with the one constructed as above.*

REMARK 1.2. Set a holomorphic 1-form $hdz := S_1 dz$ on M. In terms of the Weierstrass data (hdz, g) of M, the induced metric ds^2 and its Gaussian curvature K_{ds^2} can be rewritten respectively by

$$ds^2 = |h|^2(1 + |g|^2)|dz|^2, \qquad K_{ds^2} = -\frac{2|g'_z|^2}{|h|^2(1 + |g|^2)^3}. \tag{4}$$

Moreover, the minimal Lagrangian immersion $f\colon M \to \mathbf{C}^2$ is also given by

$$\begin{aligned} f &= \frac{1}{\sqrt{2}} e^{\mathbf{i}\,\beta/2}\left(\int ghdz - \mathbf{i}\int \overline{hdz},\ \int hdz + \mathbf{i}\int \overline{ghdz}\right) \\ &= \frac{1}{\sqrt{2}} e^{\mathbf{i}\,\beta/2}\left(-\int S_2 dz - \mathbf{i}\int \bar{S}_1 d\bar{z},\ \int \overline{\bar{S}_1 d\bar{z}} + \mathbf{i}\int \overline{S_2 dz}\right) \end{aligned} \tag{5}$$

for $(\bar{S}_1 d\bar{z}, S_2 dz) \in \Gamma(K_S^{-1} \oplus K_S)$, which is essentially same as the one proved by Hélein-Romon [11]. On the other hand, the induced metric $d\tilde{s}^2$ and its Gaussian curvature $K_{d\tilde{s}^2}$ of a minimal surface in $\mathbf{R}^3$ associated with the Weierstrass data (hdz, g) are given respectively by (cf. [26])

$$d\tilde{s}^2 = |h|^2(1 + |g|^2)^2|dz|^2, \qquad K_{d\tilde{s}^2} = -\frac{4|g'_z|^2}{|h|^2(1 + |g|^2)^4}. \tag{6}$$

With these understanding, for each $m \in \mathbf{N}$ and a Weierstrass data (hdz, g) on an open Riemann surface M, one can consider the conformal metric $ds^2_m := |h|^2(1+|g|^2)^m|dz|^2$ on M. In fact, the third author [16] has studied the precise maximal number of exceptional

values of g, provided that g is not constant and that ds_m^2 is complete. Here, we call a value that a function or map never takes an *exceptional value* of the function or map.

The purpose of the present paper is to reveal the relationship between the total curvature and the global behavior of the Gauss map of a complete minimal Lagrangian surface in $\mathbf{C}^2$. Our main theorems are Theorems 2.4 and 2.6 in Section 2. The paper is organized as follows: In Section 2, we first give a curvature bound for a minimal Lagrangian surface in $\mathbf{C}^2$ (Theorem 2.1) and show that the precise maximal number of exceptional values of the Gauss map of a complete minimal Lagrangian surface in $\mathbf{C}^2$ is "3"(Corollary 2.2). These results follow from directly Theorem 2.1 and Proposition 2.4 in [16]. Next, we give the upper bound for the number of exceptional values of the Gauss map of a complete minimal Lagrangian surface with finite total curvature in $\mathbf{C}^2$ (Theorem 2.4). In particular, we show that the precise maximal number of exceptional values of the Gauss map for this class is "2". The proof is given in Section 3.1. Moreover, by refining the Mo-Osserman argument in [23], we prove that if the Gauss map of a complete minimal Lagrangian surface in $\mathbf{C}^2$ takes on "4" distinct values only a finite number of times, then the surface has finite total curvature (Theorem 2.6). The proof is given in Section 3.2. As a corollary of the result, we obtain that if the Gauss map of a complete minimal Lagrangian surface in $\mathbf{C}^2$ which is not a Lagrangian plane omits "3" values, then it takes all other values infinitely many times (Corollary 2.7). This result is a sharping of Corollary 2.2.

Finally, the authors would like to thank Professors Katsuei Kenmotsu, Reiko Miyaoka, Masaaki Umehara and Kotaro Yamada for useful advice and continuous encouragements.

2. Main results

We first give a curvature bound for a minimal Lagrangian surface M in $\mathbf{C}^2$. Set $\omega := hdz$. Then the induced metric ds^2 on M by f is rewritten by

$$ds^2 = |h|^2(1+|g|^2)|dz|^2 = (1+|g|^2)|\omega|^2. \tag{7}$$

Applying Theorem 2.1 in [16] to the metric ds^2, we can get the following theorem.

Theorem 2.1. *Let M be a minimal Lagrangian surface in $\mathbf{C}^2$ whose Gauss map $g\colon M \to \mathbf{C}\cup\{\infty\}$ omits more than three distinct values. Then there exists a positive constant C depending on the set of exceptional values, but not M, such that for all $p \in M$ we have*

$$|K_{ds^2}(p)|^{1/2} \leq \frac{C}{d(p)},$$

where $K_{ds^2}(p)$ stands for the Gaussian curvature of M at p and $d(p)$ stands for the geodesic distance from p to the boundary of M.

Combining this with Proposition 2.4 in [16], we give the precise maximal number of exceptional values of the Gauss map of the case where M is complete.

COROLLARY 2.2. *The Gauss map of a complete minimal Lagrangian surface in* $\mathbf{C}^2$ *which is not a Lagrangian plane can omit at most three values. Moreover, let* E *be an arbitrary set of* q *points on* $\mathbf{C}\cup\{\infty\}$*, where* $q\leq 3$*. Then there exists a complete minimal Lagrangian surface in* $\mathbf{C}^2$ *whose image under the Gauss map omits precisely the set* E*.*

Note that an example of the case where $E=\emptyset$ (i.e., $q=0$) is given by the following:

$$F=(F_1,F_2)\colon \mathbf{C}\to\mathbf{C}^2,\quad (F_1,F_2)=\left(\frac{z^3}{3}+z+c_1,\,\frac{z^2}{2}+c_2\right)\quad (c_1,c_2\in\mathbf{C}).$$

Indeed, the Weierstrass data is $(hdz,g)=(zdz,(z+1)/z)$ and the resulting surface is a complete minimal Lagrangian surface in $\mathbf{C}^2$ whose Gauss map is surjective. See [17] for more details on other function-theoretic properties (e.g., ramification theorem, unicity theorem) of the Gauss map for this class.

Next, we consider the case where M is complete and has finite total curvature. From (4), the Gaussian curvature K_{ds^2} is nonpositive, and the total curvature is given by

$$\int_M K_{ds^2}dA=-\frac{1}{2}\int_M\left(\frac{2|g_z'|}{1+|g|^2}\right)^2 du\wedge dv, \tag{8}$$

where $z=u+\mathbf{i}\,v$.

PROPOSITION 2.3. *Let* M *be a complete minimal Lagrangian surface with finite total curvature in* $\mathbf{C}^2$*. Then it satisfies*

(a) *M is conformally equivalent to $\overline{M}_\gamma\backslash\{p_1,\cdots,p_k\}$, where $\overline{M}_\gamma$ is a compact Riemann surface of genus γ, and $p_1,\cdots,p_k\in\overline{M}_\gamma$ ([15]),*
(b) *The Weierstrass data (hdz,g) is extended meromorphically to $\overline{M}_\gamma$ ([26]).*

We call the points $\{p_1,\cdots,p_k\}$ *ends* of M. Thus we easily show that the total curvature of a complete minimal Lagrangian surface in $\mathbf{C}^2$ can only take the values $-2\pi m$, m a non-negative integer, or $-\infty$.

We give the upper bound for the number of exceptional values of the Gauss map of a complete minimal Lagrangian surface with finite total curvature in $\mathbf{C}^2$.

THEOREM 2.4. *Let* $M=\overline{M}_\gamma\backslash\{p_1,\cdots,p_k\}$ *be a complete minimal Lagrangian surface with finite total curvature in* $\mathbf{C}^2$ *and* $g\colon M\to\mathbf{C}\cup\{\infty\}$ *the Gauss map. Let* d *be the degree of* g *considered as a map of* $\overline{M}_\gamma$*. Assume that* M *is not a Lagrangian plane. Then we have*

$$D_g:=\sharp((\mathbf{C}\cup\{\infty\})\backslash g(M))\leq 2+\frac{2}{R},\quad \frac{1}{R}=\frac{\gamma-1+k/2}{d}<\frac{1}{2}. \tag{9}$$

In particular, the Gauss map can omit at most two values.

The number two in Theorem 2.4 is optimal. A famous example of a complete minimal Lagrangian surface with finite total curvature in $\mathbf{C}^2$ whose Gauss map omits exactly two

values is the Lagrangian catenoid ([4]). Indeed, the surface is defined as

$$F = (F_1, F_2)\colon \mathbf{C}\backslash\{0\} \to \mathbf{C}^2, \quad (F_1, F_2) = \left(z, \frac{1}{z}\right).$$

The Weierstrass data is given by $(hdz, g) = (-dz/z^2, -z^2)$ and the Gauss map omits exactly two values, 0 and ∞. Note that Castro and Urbano in [4] proved that the only Lagrangian catenoid is a minimal Lagrangian surface with circle symmetry in $\mathbf{C}^2$.

Remark 2.5. In the case of a complete minimal surface with finite total curvature in $\mathbf{R}^3$, Osserman [25] proved that the Gauss map can omit at most three values. However, at present, there exists no example whose Gauss map omits precise three values (see [6], [18] and [28] for details).

Finally, by refining the Mo-Osserman argument in [23], we get the following:

Theorem 2.6. *Let M be a complete minimal Lagrangian surface in $\mathbf{C}^2$. If the Gauss map takes on four values only a finite number of times, then M has finite total curvature.*

On the other hand, by Theorem 2.4, a complete minimal Lagrangian surface with finite total curvature whose Gauss map omits more than two values must be a Lagrangian plane. One consequence of Theorem 2.6 is therefore:

Corollary 2.7. *Let M be a complete minimal Lagrangian surface in $\mathbf{C}^2$ which is not a Lagrangian plane. If the Gauss map of M omits three values, then it takes all other values infinitely many times.*

3. Proof of Main theorems

3.1. **Proof of Theorem 2.4.** Since the total curvature is finite, we may assume that $\overline{M}_\gamma\backslash\{p_1, \cdots, p_k\}$, where $\overline{M}_\gamma$ is a compact Riemann surface of genus γ, and $p_1, \cdots, p_k \in \overline{M}_\gamma$. By a rotation of the surface, we may assume that the Gauss map g has neither zero nor pole at p_j and that the zeros and the poles of g are simple. The simple poles of g coincides with the simple zeros of hdz because the metric ds^2 is nondegenerate. By the completeness of M, hdz has poles at each end p_j ([20], [26, Lemma 9.6]). Moreover, since the surface is well-defined on M, hdz has poles of order $\mu_j \geq 2$ at p_j. Indeed, suppose that $\mu_j = 1$. Then we can expand the function $h(z)$ about each end p_j as

$$h(z) = \frac{c^j_{-1}}{z - p_j} + \sum_{n=0}^{\infty} c^j_n z^n, \quad c^j_1 \neq 0.$$

We also are able to expand the function $g(z)h(z)$ about each end p_j as

$$g(z)h(z) = \frac{d^j_{-1}}{z - p_j} + \sum_{n=0}^{\infty} d^j_n z^n$$

because g is holomorphic around p_j. By (5), if the surface is well-defined on M, then we have

$$\int_\gamma ghdz - \mathbf{i}\int_\gamma \overline{hdz} = 0, \quad \int_\gamma hdz + \mathbf{i}\int_\gamma \overline{ghdz} = 0$$

for any loop $\gamma \in H_1(M, \mathbf{Z})$. Thus we get that

$$2\pi\mathbf{i}(d^j_{-1} + \mathbf{i}\,\overline{c^j_{-1}}) = 0, \quad 2\pi\mathbf{i}(c^j_{-1} - \mathbf{i}\,\overline{d^j_{-1}}) = 0,$$

and therefore $c^j_{-1} = d^j_{-1} = 0$. Hence $\mu_j = 0$, contrary to our assumption. Applying the Riemann-Roch theorem to hdz on $\overline{M}_\gamma$, we obtain that

$$d - \sum_{j=1}^{k} \mu_j = 2\gamma - 2,$$

where d denotes the degree of g considered as a map of $\overline{M}_\gamma$. Thus we have

$$d = 2\gamma - 2 + \sum_{j=1}^{k} \mu_j \geq 2\gamma - 2 + 2k > 2\gamma - 2 + k = 2(\gamma - 1 + k/2), \tag{10}$$

and

$$\frac{1}{R} < \frac{1}{2}.$$

On the other hand, we assume that g omits D_g values. Let n_0 be the sum of the branching orders at the image of these exceptional values. Then we have

$$k \geq dD_g - n_0.$$

Let n_g be the total branching order of g on $\overline{M}_\gamma$. Then applying the Riemann-Hurwitz formula to the meromorphic function g on $\overline{M}_\gamma$, we have

$$n_g = 2(d + \gamma - 1). \tag{11}$$

Hence we have

$$D_g \leq \frac{n_0 + k}{d} \leq \frac{n_g + k}{d} = 2 + \frac{2}{R}. \tag{12}$$

□

Remark 3.1. Since

$$\int_M K_{ds^2} dA = -2\pi d,$$

the inequality in (12)

$$d \geq 2\gamma - 2 + 2k \tag{13}$$

corresponds to the Chern-Osserman inequality ([3], see also [27]) for complete minimal surfaces in $\mathbf{R}^4(= \mathbf{C}^2)$. Note that Kokubu, Umehara and Yamada [19] proved that the equality of (13) holds if and only if each end is asymptotic to a catenoid-type end or a planar end in some 3-dimensional subspace in $\mathbf{R}^4(= \mathbf{C}^2)$. In particular, all ends are embedded. Inequality (13) is useful in showing the global properties of complete minimal

Lagrangian surfaces with finite total curvature in $\mathbf{C}^2$. For instance, we can show that the following example is the unique complete minimal Lagrangian surface of the finite total curvature -2π in $\mathbf{C}^2$.

COROLLARY 3.2. *A complete minimal Lagrangian surface in* $\mathbf{C}^2$ *whose total curvature is* -2π *must have the following data:*

$$F = (F_1, F_2)\colon \mathbf{C} \to \mathbf{C}^2, \quad (F_1, F_2) = (az^2 + b,\ 2az + c) \quad (a, b, c \in \mathbf{C}). \tag{14}$$

PROOF. Since $d = 1$, $g\colon M \to S^2 (= \mathbf{C} \cup \{\infty\})$ is a conformal diffeomorphism, and M is conformally equivalent to $S^2 \setminus \{p_1, \cdots, p_k\}$. However, by (13), $2(k-1) \leq 1$. Thus $k = 1$. If $k = 1$, then M is conformally equivalent to the complex plane $\mathbf{C}$. In fact, we may identify M with $g(M) \subset \mathbf{C} \cup \{\infty\}$ and assume $(\mathbf{C} \cup \{\infty\}) \backslash g(M) = \{\infty\}$ after a suitable Möbius transformation. Then $g(z) = z$ and the holomorphic 1-form $h dz$ has no zeros on $\mathbf{C}$ because the metric ds^2 is nondegenerate. Since $h dz$ is extended meromorphically to $S^2 = \mathbf{C} \cup \{\infty\}$, $h(z)$ is a polynomial in z and therefore constant. Set that $h = 2a$ $(a \in \mathbf{C})$. Then we have

$$F_1 = \int gh\, dz = az^2 + b, \qquad F_2 = \int h dz = 2az + c.$$

□

3.2. **Proof of Theorem 2.6.** We first recall the notion of chordal distance between two distinct values in the Riemann sphere $\mathbf{C} \cup \{\infty\}$. For two distinct values α, $\beta \in \mathbf{C} \cup \{\infty\}$, we set

$$|\alpha, \beta| := \frac{|\alpha - \beta|}{\sqrt{1 + |\alpha|^2}\sqrt{1 + |\beta|^2}}$$

if $\alpha \neq \infty$ and $\beta \neq \infty$, and $|\alpha, \infty| = |\infty, \alpha| := 1/\sqrt{1 + |\alpha|^2}$. We note that, if we take v_1, $v_2 \in \mathbf{S}^2$ with $\alpha = \varpi(v_1)$ and $\beta = \varpi(v_2)$, we have that $|\alpha, \beta|$ is a half of the chordal distance between v_1 and v_2, where ϖ denotes the stereographic projection of the 2-sphere $\mathbf{S}^2$ onto $\mathbf{C} \cup \{\infty\}$.

Before proceeding to the proof of Theorem 2.6, we recall two function-theoretic lemmas.

LEMMA 3.3. [9, (8.12) in page 136] *Let g be a nonconstant meromorphic function on $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ $(0 < R \leq +\infty)$ which omits q values $\alpha_1, \cdots, \alpha_q$. If $q > 2$, then for each positive η with $\eta < (q-2)/q$, then there exists a positive constant C', depending on q and $L := \min_{i<j} |\alpha_i, \alpha_j|$, such that*

$$\frac{|g'_z|}{(1 + |g|^2) \prod_{j=1}^{q} |g, \alpha_j|^{1-\eta}} \leq C' \frac{R}{R^2 - |z|^2}. \tag{15}$$

LEMMA 3.4. [8, Lemma 1.6.7] *Let $d\sigma^2$ be a conformal flat metric on an open Riemann surface Σ. Then, for each point $p \in \Sigma$, there exists a local diffeomorphism Φ of a disk $\Delta_R = \{z \in \mathbf{C}; |z| < R\}$ $(0 < R \leq +\infty)$ onto an open neighborhood of p with $\Phi(0) = p$ such that Φ is a local isometry, that is, the pull-back $\Phi^*(d\sigma^2)$ is equal to the standard*

Euclidean metric ds^2_{Euc} *on* Δ_R *and, for a point* a_0 *with* $|a_0| = 1$, *the* Φ*-image* Γ_{a_0} *of the curve* $L_{a_0} = \{w := a_0 s; 0 < s < R\}$ *is divergent in* Σ.

Proof of Theorem 2.6. Suppose that the Gauss map g attains four distinct values $\alpha_1, \cdots, \alpha_4$ only a finite number of times. We may assume that $\alpha_4 = \infty$ after a suitable Möbius transformation. Then the assumption of the theorem implies that outside a compact subset D in M, g is holomorphic and omits three values α_1, α_2, α_3. We choose a positive number η with $0 < \eta < 1/4$ and set $\lambda := 1/(2 - 4\eta)$. Now we define a new metric

$$d\sigma^2 = |h|^{\frac{2}{1-\lambda}} \left(\frac{1}{|g'_z|} \prod_{j=1}^{3} \left(\frac{|g - \alpha_j|}{\sqrt{1 + |\alpha_j|^2}} \right)^{1-\eta} \right)^{\frac{2\lambda}{1-\lambda}} |dz|^2 \tag{16}$$

on the set $M' := \{p \in M \backslash D \,;\, g'_z(p) \neq 0\}$. Since h and g are holomorphic, $d\sigma^2$ is flat and can be smoothly extended over D. We thus obtain the metric $d\sigma^2$ on $M'' := M' \cup D$ which is flat outside the compact set D.

We prove that $d\sigma^2$ is complete on M''. This will be proved by contradiction. If $d\sigma^2$ is not complete on M'', then there exists a divergent curve $\gamma(t)$ on M'' with finite length. By removing an initial segment, if necessary, we may assume that there exists a positive distance d between the curve γ and the compact set D. Thus $\gamma \colon [0, 1) \to M'$ and, since γ is divergent on M'' with finite length, the curve $\gamma(t)$ tends to either a point where $g'_z = 0$ or else the boundary of M as $t \to 1$. However, the former case cannot occur. The reason is as follows. We assume that $\gamma(t)$ tends to a point p_0 where $g'_z = 0$ as $t \to 1$. Taking a local complex coordinate $\zeta := g'_z$ in a neighborhood of p_0 with $\zeta(p_0) = 0$, we can write

$$d\sigma^2 = |\zeta|^{-2\lambda/(1-\lambda)} w |d\zeta|^2$$

for some positive smooth function w. Since $\lambda/(1 - \lambda) > 1$, we have

$$\int_\gamma d\sigma \geq \widetilde{C} \int_\gamma \frac{|d\zeta|}{|\zeta|^{\lambda/(1-\lambda)}} = +\infty,$$

where $\widetilde{C}$ is some positive number. It contradicts that γ has finite length. We conclude that $\gamma(t)$ must tend to the boundary of M when $t \to 1$.

Then we choose a number $t_0 \in [0, 1)$ such that the length of $\gamma([t_0, 1))$ with respect to the metric $d\sigma^2$ is less than $d/3$. Since $d\sigma^2$ is flat, by Lemma 3.4, there exists an isometry Φ satisfying $\Phi(0) = \gamma(t_0)$ from a disk $\triangle_R = \{w \in \mathbf{C} \,;\, |w| < R\}$ $(0 < R \leq +\infty)$ with the standard Euclidean metric $ds^2_{\mathbf{E}}$ onto an open neighborhood of the point $\gamma(t_0)$ with the metric $d\sigma^2$, such that, for a point a_0 with $|a_0| = 1$, the Φ-image Γ_{a_0} of the line segment $L_{a_0} := \{w := a_0 s \,;\, 0 < s < R\}$ is divergent in M'. Since the length of $\gamma([t_0, 1))$ is less than $d/3$ and γ is divergent curve in M, we have $R \leq d/3$. Hence the image $\Phi(\triangle_R)$ must be bounded away from D by a distance of at least $2d/3$. Moreover, since the zeros of g'_z have been shown to be infinitely far away in the metric, Γ_{a_0} must actually go to the boundary of M.

For brevity, we denote the function $g \circ \Phi$ on $\triangle_R$ by g in the following. Since $\Phi^* d\sigma^2 = |dz|^2$, we get by (16) that

$$|h| = \left(|g_z'| \prod_{j=1}^{3} \left(\frac{\sqrt{1+|\alpha_j|^2}}{|g-\alpha_j|} \right)^{1-\eta} \right)^{\lambda}. \tag{17}$$

By Lemma 3.3, we have

$$\begin{aligned} \Phi^* ds &= |h| \sqrt{1+|g|^2} |dz| \\ &= \left(|g_z'| (1+|g|^2)^{1/2\lambda} \prod_{j=1}^{3} \left(\frac{\sqrt{1+|\alpha_j|^2}}{|g-\alpha_j|} \right)^{1-\eta} \right)^{\lambda} |dz| \\ &= \left(\frac{|g_z'|}{(1+|g|^2) \prod_{j=1}^{4} |g, \alpha_j|^{1-\eta}} \right)^{\lambda} |dz| \\ &\leq (C')^{\lambda} \left(\frac{R}{R^2-|z|^2} \right)^{\lambda} |dz|. \end{aligned}$$

Then, for the length L of Γ_{a_0} with respect to the metric ds^2, we obtain

$$L = \int_{\Gamma_{a_0}} ds = \int_{L_{a_0}} \Phi^* ds \leq (C')^{\lambda} \int_{L_{a_0}} \left(\frac{R}{R^2-|z|^2} \right)^{\lambda} |dz| \leq (C')^{\lambda} \frac{R^{1-\lambda}}{1-\lambda} < +\infty$$

because $1/2 < \lambda < 1$. In particular, the length of L is finite. We thus show that if $d\sigma^2$ were not complete on M'', then we could find a divergent curve on M with finite length in the original metric ds^2, that is, M would not be complete. However it contradicts the assumption. Hence we conclude that $d\sigma^2$ is complete on M''.

Since $d\sigma^2$ is also flat outside a compact set, the total curvature of $d\sigma^2$ is finite. Then, by the Huber theorem [15], M'' is finitely connected. We thus show that g_z' can have only a finite number of zeros and M is finitely connected. Moreover, by [24, Theorem 2.1], each annular end of M'', hence of M, is conformally equivalent to a punctured disk. Therefore the Riemann surface M must be conformally equivalent to $\overline{M} \backslash \{p_1, \cdots, p_k\}$, where $\overline{M}$ is a closed Riemann surface and $p_j \in \overline{M}$ $(j = 1, \ldots, k)$. In a neighborhood of each of p_j, g is holomorphic and omits three values. By the Picard great theorem, g cannot have an essential singularity, but must have at most a pole. Hence g extends to a meromorphic function on $\overline{M}$. Then we have

$$\int_M K_{ds^2} \, dA = -\frac{1}{2} \int_M \left(\frac{2|g_z'|}{1+|g|^2} \right)^2 du \wedge dv = -2\pi d,$$

where $z = u + \mathbf{i}v$ and d denotes the degree of g considered as a map on $\overline{M}$. In particular, M must have finite total curvature. □

References

[1] R. Aiyama, Totally real surfaces in the complex 2-space, Steps in Differential Geometry (Debrecen, 2000), pp. 15–22, Inst. Math. Inform., Debrecen, 2001.

[2] R. Aiyama, Lagrangian surfaces with circle symmetry in the complex two-space, Michigan Math. J. **52** (2004), 491–506.

[3] S. S. Chern and R. Osserman, Complete minimal surfaces in Euclidean n-space, J. Analyse Math. **19** (1967), 15–34.

[4] I. Castro and F. Urbano, On a minimal Lagrangian submanifold of $\mathbf{C}^n$ foliated by spheres, Michigan Math. J. **46** (1999), 71–82.

[5] B.-Y. Chen and J.-M. Morvan, Géométrie des surfaces lagrangiennes de $\mathbf{C}^2$, J. Math. Pures Appl. **66** (1987), 321–325.

[6] Y. Fang, On the Gauss map of complete minimal surface with finite total curvature, Indiana Univ. Math. J. **42** (1993), 1389–1411.

[7] H. Fujimoto, On the number of exceptional values of the Gauss map of minimal surfaces, J. Math. Soc. Japan **40** (1988), 237–149.

[8] H. Fujimoto, Value Distribution Theory of the Gauss Map of Minimal Surfaces in $\mathbf{R}^m$, Aspects of Mathematics, E21. Friedr. Vieweg & Sohn, Braunschweig, 1993.

[9] H. Fujimoto, Nevanlinna theory and minimal surfaces, Geometry V, 95–151, 267–272, Encyclopaedia Math. Sci., **90**, Springer, Berlin, 1997.

[10] R. Harver and H. B. Lawson Jr., Calibrated geometries, Acta Math. **148** (1982), 47–157.

[11] F. Hélein and P. Romon, Weierstrass representation of Lagrangian surfaces in four-dimensional space using spinors and quaternions, Comment. Math. Helv. **75** (2000), 668–680.

[12] F. Hélein and P. Romon, Hamiltonian stationary Lagrangian surfaces in $\mathbf{C}^2$, Comm. Anal. Geom. **10** (2002), 79–126.

[13] D. A. Hoffman and R. Osserman, The geometry of the generalized Gauss map, Mem. Amer. Math. Soc. Vol. 28, No. 236, 1980.

[14] D. A. Hoffman and R. Osserman, The Gauss map of surfaces in $\mathbf{R}^3$ and $\mathbf{R}^4$, Proc. London Math. Soc. **50** (1985), 27–56.

[15] A. Huber, On subharmonic functions and differential geometry in the large, Comment. Math. Helv. **32** (1957), 13–72.

[16] Y. Kawakami, On the maximal number of exceptional values of Gauss maps for various classes of surfaces, Math. Z. **274** (2013), 1249–1260 .

[17] Y. Kawakami, Function-theoretic properties for the Gauss maps of various classes of surfaces, preprint, arXiv:1311.1931.

[18] Y. Kawakami, R. Kobayashi and R. Miyaoka, The Gauss map of pseudo-algebraic minimal surfaces, Forum Math. **20** (2008), 1055–1069.

[19] M. Kokubu, M. Umehara and K. Yamada, Minimal surfaces that attains equality in the Chern-Osserman inequality, Contemp. Math. **308** (2002), 223–228.

[20] G. R. MacLane, On asymptotic values, abstract 603–166, Notices Amer. Math. Soc. **10** (1963), 482–483.

[21] A. Martínez, Improper affine maps, Math. Z **249** (2005), 755–766.

[22] M. J. Micallef, Stable minimal surfaces in Euclidean space, J. Differential Geom. **19** (1984), 57–84.

[23] X. Mo and R. Osserman, On the Gauss map and total curvature of complete minimal surfaces and an extension of Fujimoto's theorem, J. Differential Geom. **31** (1990), 343–355.

[24] R. Osserman, On complete minimal surfaces, Arch. Rational Mech. Anal. **13** (1963), 392–404.

[25] R. Osserman, Global properties of minimal surfaces in E^3 and E^n, Ann. of Math. **80** (1964), 340–364.

[26] R. Osserman, A Survey of Minimal Surfaces, second edition, Dover Publication Inc., New York, 1986.

[27] M. Umehara, Surfaces with singularities and Osserman-type inequalities, Proceedings of the 16th OCU International Academic Symposium "Riemann surfaces, harmonic maps and visualization", OCAMI Stud. **3**, 15–28, 2009.

[28] A. Weitsman and F. Xavier, Some function theoretic properties of the Gauss map for hyperbolic complete minimal surfaces, Michigan Math. J. **34** (1987), 275–283.

Institute of Mathematics,
University of Tsukuba,
Tsukuba 305-8571, Japan
E-mail address: aiyama@math.tsukuba.ac.jp

Department of Mathematics,
Tokyo Institute of Technology,
Tokyo 152-8551, Japan
E-mail address: akutagawa@math.titech.ac.jp

Graduate School of Natural Science and Technology,
Kanazawa university,
Kanazawa, 920-1192, Japan
E-mail address: y-kwkami@se.kanazawa-u.ac.jp